
# NC Violent Death Reporting System

## YOUTH SUICIDE IN NORTH CAROLINA, AGES 10-24, 2013 - 2022

The North Carolina Violent Death Reporting System (NC-VDRS) is a CDC-funded statewide surveillance system that collects detailed information on violent deaths that occur in North Carolina including: homicide, suicide, unintentional firearm deaths, legal intervention, and deaths of undetermined intent. NC-VDRS uses information from death certificates, medical examiner reports, and law enforcement reports to understand and aid in the prevention of violent deaths. For more information, please visit https://www.injuryfreenc.ncdhhs.gov/About/ncVDRS.htm.

## Youth Suicide in North Carolina, by Sex and Age: NC-VDRS, 2013 - 2022



- From 2013 to 2022, 3,975 North Carolina residents ages 10 to 24 died as a result of violence. Of these violent deaths, 1,908 (48.0%) were suicides.
- Males consistently had a higher number of suicides than females, regardless of age.
- The number of suicide deaths peaked for males at age 22 (214 suicides) and for females at age 23 (51 suicides).
- Sixty-six percent of all youth suicide victims were identified as non-Hispanic (NH) White, 19.8% as NH Black, 9.1% as Hispanic, and 5.6% as belonging to other racial/ethnic groups.


- Overall, the most common method of suicide among youth was firearms (55.4%), followed by hanging, strangulation, suffocation (33.6%) and poisoning (5.8%).
- Only 5.2% of suicides involved a method other than firearms, hanging, strangulation, suffocation, or poisoning.

# If you or someone you know needs support now,




call or text 988 or chat 988lifeline.org

## Method of Youth Suicides: NC-VDRS, 2013 - 2022



\*Other includes falls, drowning, motor vehicle, fire/burns and other causes of suicide.

# Precipitating Circumstances of Youth Suicide by Sex: NC-VDRS, 2013 - 2022



- Sixty-one percent (60.5%) of female and 46.1% of male suicide victims were characterized as having a current mental health problem when they died by suicide.
- A similar trend was seen for mental health treatment. Females (45.9%) were more likely than males (29.7%) to be receiving treatment for a current mental health problem at the time of suicide.
- Thirty percent (30.1%) of female and 28.0% of male suicide victims had an intimate partner problem.
- Forty-six percent (45.9%) of female and 46.3% of male suicide victims had experienced a recent or imminent crisis.
- Twenty-nine percent (29.0%) of female and 26.2% of male suicide victims had disclosed their suicide intent to someone else before they died.
- Females (40.1%) were more likely to leave a suicide note than males (28.0%).

More infromation on suicide prevention efforts can be found at:

#### **State Resource Partners**

Division of Mental Health,

#### **Developmental Disabilities and Substance Use Services**

https://www.ncdhhs.gov/divisions/mhddsas

#### Office of the Chief Medical Examiner

https://www.ocme.dhhs.nc.gov/

#### Injury and Violence Prevention's Suicide Prevention Website

https://injuryfreenc.dph.ncdhhs.gov/preventionResources/Suicide.htm

#### **Division of Child and Family Wellbeing**

https://www.ncdhhs.gov/divisions/division-child-and-family-well-being

<u>National Resources</u> The Suicide Prevention Resource Center

http://www.sprc.org/

The American Foundation for Suicide Prevention

https://afsp.org/

The National Suicide Prevention Lifeline

call or text 988

or chat 988lifeline.org

The North Carolina Violent Death Reporting System is supported by Cooperative Agreement 5 NU17CE924955-02-00 from the Centers for Disease Control and Prevention (CDC).





NCDHHS Division of Public Health / www.publichealth.nc.gov / Injury Epidemiology & Surveillance Unit/ 919-707-5425
NC Violent Death Reporting System / 919-707-5432

State of North Carolina / Department of Health and Human Services / www.ncdhhs.gov NCDHHS is an equal opportunity employer and provider. 2022 FINAL DATA 7/11/2024

Please see the NC-VDRS 2022 Annual Report for additional data and technical information:

https://www.injuryfreenc.ncdhhs.gov/DataSurveillance/ViolentDeathData.htm

<sup>\*</sup>Circumstances are known for 92% of males (n=1413) and 96% of females (n=362)